

ON THE DISTRIBUTION OF LATTICE POINTS IN THIN ANNULI

C.P. HUGHES AND Z. RUDNICK

ABSTRACT. We show that the number of lattice points lying in a thin annulus has a Gaussian value distribution if the width of the annulus tends to zero sufficiently slowly as we increase the inner radius.

1. INTRODUCTION

Let $N(t)$ be the number of integer lattice points in a disk of radius t centered at the origin. Thus $N(t) = \sum_{n \leq t^2} r(n)$ where $r(n)$ is the number of ways of writing $n = x^2 + y^2$ as a sum of two squares. As is well known, $N(t)$ is asymptotic to the area πt^2 of the disk. Much effort has gone into understanding the growth of the remainder term. Heath-Brown [9] considered the *distribution* of the normalized remainder term $(N(t) - \pi t^2)/\sqrt{t}$, and proved that it has a limiting value distribution in the sense that there exists a probability distribution function ν such that for any interval $\mathcal{A}$,

$$\frac{1}{T} \text{meas} \left\{ t \in [T, 2T] : \frac{N(t) - \pi t^2}{\sqrt{t}} \in \mathcal{A} \right\} \rightarrow \int_{\mathcal{A}} \nu(x) \, dx$$

where the measure is the ordinary Lebesgue measure. It is known that $\nu(x)$ is not the Gaussian measure, for instance the tails have been shown to decay roughly like $\exp(-x^4)$, [4, 10].

Bleher, Dyson and Lebowitz [5, 6, 3] investigated the distribution of a similarly scaled remainder term of the number $N(t, \rho) := N(t + \rho) - N(t)$ of lattice points in an annulus of inner radius t and width $\rho(t)$ depending on t . The “expected” number of points is the area $\pi(2t\rho + \rho^2)$ of the annulus. Define a normalized remainder term by

$$S(t, \rho) := \frac{N(t + \rho) - N(t) - \pi(2t\rho + \rho^2)}{\sqrt{t}}.$$

The picture that emerges is that there is a number of distinct regimes:

- The “global”, or “macroscopic”, regime $\rho(t) \rightarrow \infty$ (but $\rho = o(t)$), in which case Bleher and Lebowitz [6] show that $S(t, \rho)$ has a limiting distribution with tails which decay roughly as $\exp(-x^4)$. In fact the distribution is that of the difference of two i.i.d. random variables whose distribution is the limiting distribution of $(N(t) - \pi t^2)/\sqrt{t}$.
- The intermediate, or “mesoscopic”, regime $\rho \rightarrow 0$ but $\rho t \rightarrow \infty$. The variance of $S(t, \rho)$ is given by [7]

$$\frac{1}{T} \int_T^{2T} |S(t, \rho)|^2 dt \sim \sigma^2 := 16\rho \log \frac{1}{\rho}$$

Date: 12 September 2003.

and Bleher and Lebowitz [6] conjectured that $S(t, \rho)/\sigma$ has a standard Gaussian distribution.

- The “saturation” regime $0 < \rho(t) < \infty$ is fixed as $t \rightarrow \infty$, where it has been shown [6] that $S(t, \rho)$ has a distribution with rapidly decaying tails. As $\rho \rightarrow \infty$ the distribution converges to that found in the macroscopic regime, and as $\rho \rightarrow 0$ it converges to the conjectured mesoscopic distribution.
- The local regime, $\rho \approx 1/t$. If the annulus were centered at a generic point rather than at a lattice point, or if we consider “generic” lattices instead of the integer lattice $\mathbb{Z}^2$, then it is consistent with conjectures of Berry and Tabor [1] that the statistics are Poissonian; see [15, 8, 12] for some progress on this, as well as [16, 11, 13].

In this paper we prove part of the Gaussian distribution conjecture of Bleher and Lebowitz. We will show that $S(t, \rho)$ has a Gaussian distribution when ρ shrinks to zero sufficiently slowly:

Theorem 1. *If $\rho \rightarrow 0$ but $\rho \gg T^{-\delta}$ for all $\delta > 0$, then for any interval $\mathcal{A}$*

$$\lim_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ t \in [T, 2T] : \frac{S(t, \rho)}{\sigma} \in \mathcal{A} \right\} = \frac{1}{\sqrt{2\pi}} \int_{\mathcal{A}} e^{-x^2/2} dx$$

where $\sigma^2 = 16\rho \log \frac{1}{\rho}$.

The structure of the argument is as follows: We replace the sharp counting function $N(t)$ by a smooth counting function $\tilde{N}_M(t)$ whose smoothness parameter $M = M(T)$ depends on T (note that though t and T are formally independent, we always think of t as being around T). Since we are only interested in $\rho \rightarrow 0$ we will set $\rho = 1/L$ where $L = L(T)$ tends to infinity with T , and we define the corresponding normalized remainder term to be

$$\tilde{S}_{M,L}(t) := \frac{\tilde{N}_M(t + 1/L) - \tilde{N}_M(t) - 2\pi/L - \pi/L^2}{\sqrt{t}}$$

We compute the moments of $\tilde{S}_{M,L}(t)$ when t is chosen at random with respect to a smooth measure. We show in Section 3 that the m^{th} moment of $\tilde{S}_{M,L}/\sigma$ converges to that of a standard normal random variable provided $L \ll T^{\nu(m)}$, with $0 < \nu(m) < 1/(2^{m-1} - 1)$. Thus $\tilde{S}_{M,L}$ has a normal distribution if $L \rightarrow \infty$ but $L \ll T^\delta$ for all $\delta > 0$. In Section 4 we show that the variance of the difference $(S(t, 1/L) - \tilde{S}_{M,L}(t))/\sigma$ goes to zero, and hence $S(t, \rho)/\sigma$ has a normal distribution with respect to the smooth measure. Finally we use an approximation argument to pass from smooth measures to the Lebesgue measure used in Theorem 1.

2. SMOOTHING

To obtain Theorem 1 we will replace sharp cutoffs by smooth ones. First, we will replace Lebesgue measure with a smooth average of t around T , that is we pick t at random by taking a smooth function $\omega \geq 0$, of total mass unity, such that both ω and its Fourier transform $\hat{\omega}$ are rapidly decaying, in the sense that for any $A > 2$,

$$(1) \quad \omega(t) \ll \frac{1}{(1 + |t|)^A}, \quad \hat{\omega}(t) \ll \frac{1}{(1 + |t|)^A}$$

for all t . (In fact we also choose ω to be supported on the positive reals as this makes the analysis simpler).

Define the averaging operator

$$(2) \quad \langle f \rangle = \frac{1}{T} \int_{-\infty}^{\infty} f(t) \omega\left(\frac{t}{T}\right) dt$$

(this is the expected value of f with respect to this measure), and let $\mathbb{P}_{\omega,T}$ be the associated probability measure:

$$\mathbb{P}_{\omega,T}(f \in \mathcal{A}) = \frac{1}{T} \int_{-\infty}^{\infty} \mathbb{1}_{\mathcal{A}}(f(t)) \omega\left(\frac{t}{T}\right) dt$$

(Throughout the paper we will extend $N(t)$, $S(t, \rho)$ and similar functions, initially defined for $t > 0$, to the whole real line. Since $\omega(t) = 0$ for $t \leq 0$ we are free to choose whichever extension makes the analysis most simple).

We will also smooth the edges of the circle, and show that this modified counting function has a Gaussian distribution: Let χ be the indicator function of the unit disc, and ψ a smooth, even, function on the real line, of total mass unity, whose Fourier transform $\widehat{\psi}$ is smooth and has compact support. Define a rotationally symmetric function Ψ on $\mathbb{R}^2$ by setting $\widehat{\Psi}(\vec{y}) = \widehat{\psi}(|\vec{y}|)$ where $|\vec{y}|$ denotes the standard Euclidean norm of $\vec{y} \in \mathbb{R}^2$, and where the Fourier transform is

$$\widehat{f}(\vec{y}) = \int_{\mathbb{R}^2} f(\vec{x}) e^{-2\pi i \langle \vec{x}, \vec{y} \rangle} d\vec{x}$$

with $\langle \vec{x}, \vec{y} \rangle$ the usual Euclidean inner product. For $\epsilon > 0$ set

$$(3) \quad \Psi_{\epsilon}(\vec{x}) = \frac{1}{\epsilon^2} \Psi\left(\frac{\vec{x}}{\epsilon}\right)$$

Now set $\chi_{\epsilon} = \chi * \Psi_{\epsilon}$ to be the convolution of χ and Ψ_{ϵ} , which is a smoothed indicator function of the unit disc with “fuzziness” of width ϵ , in the sense that $0 \leq \chi_{\epsilon} \leq 1$, and if ψ (rather than its Fourier transform $\widehat{\psi}$) had compact support then $\chi - \chi_{\epsilon}$ would be concentrated in the shell $1 - \epsilon < |\vec{x}| < 1 + \epsilon$. Due to the rapid decay of tails, this is essentially still the case when ψ is in the Schwarz class, as it is for us.

Now take $\epsilon = 1/t\sqrt{M}$ where $M = M(T)$ depends on T and tends to infinity with T , and define a smooth counting function, or smooth linear statistic, by

$$\widetilde{N}_M(t) = \sum_{\vec{n} \in \mathbb{Z}^2} \chi_{\epsilon}\left(\frac{\vec{n}}{t}\right)$$

This counts lattice points in a “fuzzy circle” of radius about t , with fuzziness about $t\epsilon = 1/\sqrt{M}$.

The number of lattice points in a smooth annulus of inner radius t and width ρ is therefore given by $\widetilde{N}_M(t + \rho) - \widetilde{N}_M(t)$. Since we are interested in radii t in an interval $[T, 2T]$, we will in what follows freeze the width of the annulus to be $\rho(T)$ as t varies in $[T, 2T]$ rather than allowing it to vary with t ; this will simplify some of the calculations. Furthermore, since from henceforth we are only concerned with $\rho \rightarrow 0$, we will set $\rho = 1/L$, and let $L(T) \rightarrow \infty$ as $T \rightarrow \infty$.

Set

$$(4) \quad \widetilde{S}_{M,L} = \frac{\widetilde{N}_M(t + 1/L) - \widetilde{N}_M(t) - 2\pi t/L - \pi/L^2}{\sqrt{t}}$$

The width of the smoothed sides of $\tilde{N}_M$ is $\mathcal{O}(\epsilon t) = \mathcal{O}(1/\sqrt{M})$. In order for $\tilde{S}_{M,L}$ to approximate $S(t, 1/L)$, it must be that $1/L$ is much larger than the width of the sides, so we insist that $L/\sqrt{M} \rightarrow 0$.

We will show:

Theorem 2. *Suppose that $M(T)$, $L(T)$ are increasing to infinity with T such that $M = \mathcal{O}(T^\delta)$ for all $\delta > 0$, and $L/\sqrt{M} \rightarrow 0$, then for any interval $\mathcal{A}$,*

$$\lim_{T \rightarrow \infty} \mathbb{P}_{\omega, T} \left\{ \frac{\tilde{S}_{M,L}}{\sigma} \in \mathcal{A} \right\} = \frac{1}{\sqrt{2\pi}} \int_{\mathcal{A}} e^{-x^2/2} dx$$

where $\tilde{S}_{M,L}$ is given by (4) and

$$\sigma^2 = \frac{16 \log L}{L}$$

Remark. The arguments given below for the proof of Theorem 2 will also prove a central limit theorem for smooth linear statistics in higher dimensions: Defining $\chi_\epsilon = \chi * \Psi_\epsilon$ where χ is the indicator function of the unit ball and Ψ_ϵ is defined in analogy with (3), we have a smooth counting function $\tilde{N}_M(t) := \sum_{\vec{n} \in \mathbb{Z}^d} \chi_\epsilon(\frac{\vec{n}}{t})$ where as before $\epsilon = 1/t\sqrt{M}$.

The asymptotic behaviour of $\tilde{N}_M(t)$ is given by $c_d t^d$, with c_d the volume of the unit ball in $\mathbb{R}^d$.

It may then be shown that if $M = \mathcal{O}(T^\delta)$ for all $\delta > 0$, then the distribution of the normalized remainder term $\tilde{S}_M(t) = \frac{\tilde{N}_M(t) - c_d t^d}{t^{(d-1)/2}}$ when averaged over t around T weakly converges to a Gaussian with mean zero and variance

$$\sigma^2 = \begin{cases} \frac{2}{\pi^2} K_3 \log M & \text{when } d = 3 \\ \frac{d-1}{\pi^2} K_d \int_0^\infty y^{d-4} \hat{\psi}(y)^2 dy M^{(d-3)/2} & \text{when } d \geq 4 \end{cases}$$

where

$$K_d = \frac{4^{d-1} \pi^{d-1/2}}{2^d - 1} \frac{\Gamma(\frac{1}{2}d - \frac{1}{2})}{\Gamma(d) \Gamma(\frac{1}{2}d)} \frac{\zeta(d-1)}{\zeta(d)}.$$

3. THE DISTRIBUTION OF $\tilde{N}_M$

Lemma 3. *As $t \rightarrow \infty$, we have*

$$\tilde{N}_M(t) = \pi t^2 - \frac{\sqrt{t}}{\pi} \sum_{n=1}^\infty \frac{r(n)}{n^{3/4}} \cos(2\pi t \sqrt{n} + \frac{1}{4}\pi) \hat{\psi}(\sqrt{\frac{n}{M}}) + \mathcal{O}\left(\frac{1}{\sqrt{t}}\right)$$

with the error term independent of M .

Proof. By Poisson summation,

$$\begin{aligned} \tilde{N}_M(t) &:= \sum_{\vec{n} \in \mathbb{Z}^2} (\chi * \Psi_\epsilon) \left(\frac{\vec{n}}{t} \right) \\ (5) \quad &= t^2 \sum_{\vec{k} \in \mathbb{Z}^2} \hat{\chi}(t\vec{k}) \hat{\Psi}_\epsilon(t\vec{k}) \end{aligned}$$

Changing into polar coordinates, and using the fact that χ is rotationally symmetric, the 2-dimensional Fourier transform of χ is

$$\begin{aligned}\widehat{\chi}(\vec{y}) &= \int_0^1 r \int_0^{2\pi} e^{-2\pi i r |\vec{y}| \cos \theta} d\theta dr \\ &= \frac{-\cos(2\pi|\vec{y}| + \frac{1}{4}\pi)}{\pi|\vec{y}|^{3/2}} + \mathcal{O}\left(\frac{1}{|\vec{y}|^{5/2}}\right)\end{aligned}$$

as $|\vec{y}| \rightarrow \infty$. By its definition in (3), $\widehat{\Psi}_\epsilon(\vec{y}) = \widehat{\Psi}(\epsilon\vec{y}) = \widehat{\psi}(\epsilon|\vec{y}|)$. Therefore, inserting this into (5), treating the mean (when $\vec{k} = \vec{0}$) separately, and setting $\epsilon = 1/t\sqrt{M}$,

$$\begin{aligned}\widetilde{N}_M(t) &= \pi t^2 - \frac{\sqrt{t}}{\pi} \sum_{\vec{k} \neq \vec{0}} \left\{ \frac{\cos(2\pi t|\vec{k}| + \frac{1}{4}\pi)}{|\vec{k}|^{3/2}} \widehat{\psi}(\epsilon t|\vec{k}|) + \mathcal{O}\left(\frac{1}{t} \frac{\widehat{\psi}(\epsilon t|\vec{k}|)}{|\vec{k}|^{5/2}}\right) \right\} \\ &= \pi t^2 - \frac{\sqrt{t}}{\pi} \sum_{n=1}^{\infty} \frac{r(n)}{n^{3/4}} \cos(2\pi t\sqrt{n} + \frac{1}{4}\pi) \widehat{\psi}(\sqrt{\frac{n}{M}}) + \mathcal{O}\left(\frac{1}{\sqrt{t}}\right)\end{aligned}$$

with the constant implicit in the error term independent of $M(T)$. $\square$

Note that the compact support of $\widehat{\psi}$ means that the sum truncates at $n \approx M$. Thus we need $M \gg 1$ in order for there to be any terms in the sum.

Now, since

$$\widetilde{S}_{M,L} = \frac{\widetilde{N}_M(t + 1/L) - \widetilde{N}_M(t) - \pi(2t/L + 1/L^2)}{\sqrt{t}}$$

then for $t \geq 1$ and $L \geq 1$,

$$\begin{aligned}\widetilde{S}_{M,L} &= \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{r(n)}{n^{3/4}} \left[\cos(2\pi t\sqrt{n} + \frac{\pi}{4}) - \cos(2\pi(t + \frac{1}{L})\sqrt{n} + \frac{\pi}{4}) \right] \widehat{\psi}(\sqrt{\frac{n}{M}}) + \mathcal{O}\left(\frac{1}{t}\right) \\ (6) \quad &= \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{r(n)}{n^{3/4}} \sin\left(\frac{\pi\sqrt{n}}{L}\right) \sin\left(2\pi(t + \frac{1}{2L})\sqrt{n} + \frac{\pi}{4}\right) \widehat{\psi}(\sqrt{\frac{n}{M}}) + \mathcal{O}\left(\frac{1}{t}\right)\end{aligned}$$

Note that we have three independent variables. The variable t , which we always consider to be large, is the radius of the annulus. This is the variable we average over. The width of the annulus is $1/L$. Since we want a thin annulus, $L \rightarrow \infty$, and Gaussian behaviour is not seen if this condition does not hold. The annulus does not have sharp sides, but smoothed edges, and the third independent variable is M ; the larger M is, the sharper the annulus' sides (in the sense that it better approximates the indicator function). We must have $L/\sqrt{M} \rightarrow 0$ in order for the annulus to have some width, and not be "just sides". That is, the annulus shouldn't be too smooth.

Proof of Theorem 2. First we show the mean is $\mathcal{O}(1/T)$. Since $\omega(t)$ is real,

$$\langle \sin(2\pi(t + \frac{1}{2L})\sqrt{n} + \frac{1}{4}\pi) \rangle = \Im \left\{ \widehat{\omega}(-T\sqrt{n}) e^{i\pi(\frac{\sqrt{n}}{L} + \frac{1}{4})} \right\} \ll \frac{1}{T^A n^{A/2}}$$

for any $A > 2$, where we have used the rapid decay of $\widehat{\omega}$. Thus

$$\begin{aligned} \langle \widetilde{S}_{M,L} \rangle &\ll \sum_{n=1}^{\infty} \frac{r(n)}{n^{3/4}} \frac{1}{T^A n^{A/2}} + \mathcal{O}\left(\frac{1}{T}\right) \\ &= \mathcal{O}\left(\frac{1}{T}\right) \end{aligned}$$

Setting

$$(7) \quad \mathcal{M}_m := \left\langle \left(\frac{2}{\pi} \sum_{n=1}^{\infty} \frac{r(n)}{n^{3/4}} \sin\left(\frac{\pi\sqrt{n}}{L}\right) \sin\left(2\pi\left(t + \frac{1}{2L}\right)\sqrt{n} + \frac{1}{4}\pi\right) \widehat{\psi}\left(\sqrt{\frac{n}{M}}\right) \right)^m \right\rangle$$

then from (6), the Cauchy-Schwartz inequality implies that the m^{th} moment of $\widetilde{S}_{M,L}$ is

$$\begin{aligned} (8) \quad \langle (\widetilde{S}_{M,L})^m \rangle &= \left\langle \left\{ \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{r(n)}{n^{3/4}} \sin\left(\frac{\pi\sqrt{n}}{L}\right) \sin\left(2\pi\left(t + \frac{1}{2L}\right)\sqrt{n} + \frac{1}{4}\pi\right) \widehat{\psi}\left(\sqrt{\frac{n}{M}}\right) + \mathcal{O}\left(\frac{1}{T}\right) \right\}^m \right\rangle \\ &= \mathcal{M}_m + \mathcal{O}\left(\sum_{j=1}^m \binom{m}{j} \frac{\sqrt{\mathcal{M}_{2m-2j}}}{T^j}\right) \end{aligned}$$

The conditions of Theorem 2 are that $M = \mathcal{O}(T^\delta)$ for all $\delta > 0$, and that $L \rightarrow \infty$ in such a way that $L/\sqrt{M} \rightarrow 0$. In which case, Proposition 4 allows us to deduce that $\sigma^2 := \mathcal{M}_2 \sim \frac{16 \log L}{L}$, and Proposition 5 shows that for all $m > 2$,

$$\frac{\mathcal{M}_m}{\sigma^m} = \begin{cases} \frac{m!}{2^{m/2}(m/2)!} + \mathcal{O}\left(\frac{1}{L^{1-\delta'}}\right) & \text{if } m \text{ is even} \\ \mathcal{O}\left(\frac{1}{L^{1-\delta'}}\right) & \text{if } m \text{ is odd} \end{cases}$$

These are the moments of the standard normal distribution, and inserting these into (8), we see this is sufficient to prove that the distribution of $\widetilde{S}_{M,L}/\sigma$ weakly converges as $T \rightarrow \infty$ to a Gaussian with mean zero and variance 1. $\square$

3.1. The variance.

Proposition 4. *If $M = \mathcal{O}(T^{2(1-\delta)})$ for fixed $\delta > 0$, then the variance of $\widetilde{S}_{M,L}$ is asymptotic to*

$$(9) \quad \sigma^2 := \frac{2}{\pi^2} \sum_{n=1}^{\infty} \frac{r(n)^2}{n^{3/2}} \sin^2\left(\frac{\pi\sqrt{n}}{L}\right) \widehat{\psi}^2\left(\sqrt{\frac{n}{M}}\right)$$

If $L \rightarrow \infty$ but $L/\sqrt{M} \rightarrow 0$, then

$$(10) \quad \sigma^2 \sim \frac{16 \log L}{L}$$

Proof. Expanding out (7) we have

$$\begin{aligned} \mathcal{M}_2 &= \frac{4}{\pi^2} \sum_{m,n} \frac{r(m)r(n) \sin(\pi\sqrt{m}/L) \sin(\pi\sqrt{n}/L) \widehat{\psi}(\sqrt{\frac{m}{M}}) \widehat{\psi}(\sqrt{\frac{n}{M}})}{(mn)^{3/4}} \times \\ &\quad \times \langle \sin\left(2\pi\left(t + \frac{1}{2L}\right)\sqrt{m} + \frac{1}{4}\pi\right) \sin\left(2\pi\left(t + \frac{1}{2L}\right)\sqrt{n} + \frac{1}{4}\pi\right) \rangle \end{aligned}$$

Now the average on the bottom line is

$$\begin{aligned} & \frac{1}{4} \left[\widehat{\omega} \left(T(\sqrt{m} - \sqrt{n}) \right) e^{i\pi \frac{1}{L}(\sqrt{n} - \sqrt{m})} + \widehat{\omega} \left(T(\sqrt{n} - \sqrt{m}) \right) e^{i\pi \frac{1}{L}(\sqrt{m} - \sqrt{n})} \right. \\ & \left. - \widehat{\omega} \left(T(\sqrt{m} + \sqrt{n}) \right) e^{-i\pi(\frac{1}{2} + \frac{1}{L}(\sqrt{m} + \sqrt{n}))} - \widehat{\omega} \left(-T(\sqrt{m} + \sqrt{n}) \right) e^{i\pi(\frac{1}{2} + \frac{1}{L}(\sqrt{m} + \sqrt{n}))} \right] \end{aligned}$$

The support condition on $\widehat{\psi}$ means that m, n are both constrained to be $\mathcal{O}(M)$, and so either $m = n$ or else $|\sqrt{m} - \sqrt{n}| \gg \frac{1}{\sqrt{M}}$. Using the bound $\widehat{\omega}(t) \ll (1 + |t|)^{-A}$ for all $A > 0$, the off-diagonal terms contribute at most

$$\sum_{1 \leq n \neq m \leq M} \left(\frac{\sqrt{M}}{T} \right)^A \ll \frac{M^{A/2+2}}{T^A} \ll T^{4-\delta A}$$

using the assumption $M = \mathcal{O}(T^{2(1-\delta)})$. Therefore, for any $B > 0$,

$$\mathcal{M}_2 = \frac{2}{\pi^2} \sum_{n=1}^{\infty} \frac{r(n)^2}{n^{3/2}} \sin^2 \left(\frac{\pi \sqrt{n}}{L} \right) \widehat{\psi}^2 \left(\sqrt{\frac{n}{M}} \right) + \mathcal{O}(T^{-B})$$

Define σ^2 to be the infinite sum above. Since $r(n) \ll n^\eta$ for all $\eta > 0$, σ^2 is bounded for all L . To find the asymptotics as $L \rightarrow \infty$, we use a formula of Ramanujan [14]¹:

$$\sum_{n \leq X} r(n)^2 = 4X \log X + O(X).$$

We then have

$$\begin{aligned} \sigma^2 &:= \frac{2}{\pi^2} \sum_{n=1}^{\infty} \frac{r(n)^2}{n^{3/2}} \sin^2 \left(\frac{\pi \sqrt{n}}{L} \right) \widehat{\psi}^2 \left(\sqrt{\frac{n}{M}} \right) \\ &\sim \frac{8}{\pi^2} \int_1^{\infty} \frac{\log x}{x^{3/2}} \sin^2 \left(\frac{\pi \sqrt{x}}{L} \right) \widehat{\psi}^2 \left(\sqrt{\frac{x}{M}} \right) dx \\ &= \frac{32}{L\pi^2} \int_{1/L}^{\infty} \log(yL) \frac{\sin^2(\pi y)}{y^2} \widehat{\psi}^2 \left(\frac{yL}{\sqrt{M}} \right) dy \\ &\sim \frac{\log L}{L} \frac{32}{\pi^2} \int_0^{\infty} \frac{\sin^2(\pi y)}{y^2} \widehat{\psi}^2 \left(\frac{yL}{\sqrt{M}} \right) dy \end{aligned}$$

on changing variables to $x = y^2 L^2$, and using the fact that we assume that $L \rightarrow \infty$. Now using the additional restriction (caused by the fuzziness of the annulus' sides) that $L/\sqrt{M} \rightarrow 0$, we see that since $\widehat{\psi}(yL/\sqrt{M}) \sim 1$ for all $y = o(\sqrt{M}/L)$, the integral can be evaluated asymptotically to equal $\pi^2/2$, and so

$$\sigma^2 \sim \frac{16 \log(L)}{L}$$

Since $L = o(T^{1-\delta})$, the error terms in (8) are all smaller than σ^2 , and so the variance of $\widetilde{S}_{M,L}$ is asymptotic to σ^2 as $T \rightarrow \infty$. $\square$

The constraints on M , that $M = \mathcal{O}(T^{2-2\delta})$ but $L/\sqrt{M} \rightarrow 0$, illustrate the role of smoothing. The first constraint, that M is not too big, comes from requiring that the annulus is sufficiently smooth to easily handle the averages (to enable us to reduce to the diagonal). The second constraint, that M is not too small, is to ensure the function is not too smooth, so that the width of the edges is greater

¹We wish to thank Bob Vaughan for pointing this reference out to us.

than the size of the annulus. (That $L \rightarrow \infty$ forces M to go to infinity. If it did not, the function would be so smooth as to have no fluctuations!)

3.2. The higher moments.

Proposition 5. *For fixed $\delta > 0$, if $M = \mathcal{O}\left(T^{2(1-\delta)/(2^{m-1}-1)}\right)$, and if $L \rightarrow \infty$ such that $L/\sqrt{M} \rightarrow 0$, then for arbitrary $\delta' > 0$,*

$$\frac{\mathcal{M}_m}{\sigma^m} = \begin{cases} \frac{m!}{2^{m/2}(m/2)!} + \mathcal{O}\left(\frac{1}{L^{1-\delta'}}\right) & \text{if } m \text{ is even} \\ \mathcal{O}\left(\frac{1}{L^{1-\delta'}}\right) & \text{if } m \text{ is odd} \end{cases}$$

where $\mathcal{M}_m$ is given in (7) and σ^2 is given in (9).

We will need to give lower bounds for alternating sums $\sum \pm \sqrt{n_j}$. To do so, we use the following lemma, a form of Liouville's theorem, (cf [9]):

Lemma 6. *For $j = 1, \dots, m$, let $n_j \leq M$ be positive integers. Then*

- *either $\sum \epsilon_j \sqrt{n_j} = 0$ for some $\epsilon_j = \pm 1$,*
- *or for all $\epsilon_j = \pm 1$,*

$$\left| \sum_{j=1}^m \epsilon_j \sqrt{n_j} \right| \geq \frac{1}{(m\sqrt{M})^{2^{m-1}-1}}$$

Proof. Assume that $\sum \epsilon_j \sqrt{n_j} \neq 0$ for all choices of $\epsilon_j = \pm 1$. Then

$$P := \prod_{\epsilon_j = \pm 1} \left(\sum_{j=1}^m \epsilon_j \sqrt{n_j} \right)$$

is non-zero. By Galois theory, since $\sum \epsilon_j \sqrt{n_j}$ is an algebraic number, and P is the product over all possible symmetries, P is an integer. Since we assumed that no term in P vanishes, $|P| \geq 1$. Since both $\sum \epsilon_j \sqrt{n_j}$ and $-\sum \epsilon_j \sqrt{n_j}$ are terms in P , if

$$Q := \prod_{\substack{\epsilon_j = \pm 1 \\ j=2,3,\dots,m}} \left(\sqrt{n_1} + \sum_{j=2}^m \epsilon_j \sqrt{n_j} \right)$$

then $P = (-1)^{2^{m-1}-1} Q^2$, and so $|Q| = \sqrt{|P|} \geq 1$.

By assumption $n_j \leq M$ for all j , and so, independent of the ϵ_j ,

$$\left| \sqrt{n_1} + \sum_{j=2}^m \epsilon_j \sqrt{n_j} \right| \leq m\sqrt{M}$$

and so for any $\eta_j = \pm 1$,

$$\left| \sqrt{n_1} + \sum_{j=2}^m \eta_j \sqrt{n_j} \right| = \frac{|Q|}{\prod^* \left| \sqrt{n_1} + \sum_{j=2}^m \epsilon_j \sqrt{n_j} \right|} \geq \frac{1}{(m\sqrt{M})^{2^{m-1}-1}}$$

where $\prod^*$ denotes the product over all ϵ_j distinct from η_j , there being $2^{m-1} - 1$ terms in such a product. $\square$

From this, it is simple to derive the following lemma.

Lemma 7. For $j = 1, \dots, m$ let $n_j \leq M$ be positive integers, and let $\epsilon_j = \pm 1$ be such that

$$\sum_{j=1}^m \epsilon_j \sqrt{n_j} \neq 0$$

Then

$$\left| \sum_{j=1}^m \epsilon_j \sqrt{n_j} \right| \geq \frac{1}{(m\sqrt{M})^{2^{m-1}-1}}$$

Proof. Either $\sum \eta_j \sqrt{n_j} \neq 0$ for any choice of $\eta_j = \pm 1$, and we are done by Lemma 6, or else there exists a (strict) subset $S \subsetneq \{1, \dots, m\}$ such that

$$\sum_{j \in S} \epsilon_j \sqrt{n_j} - \sum_{j \notin S} \epsilon_j \sqrt{n_j} = 0$$

so that

$$\left| \sum_{j=1}^m \epsilon_j \sqrt{n_j} \right| = 2 \left| \sum_{j \in S} \epsilon_j \sqrt{n_j} \right|$$

Note that, by assumption, $\sum_{j \in S} \epsilon_j \sqrt{n_j} \neq 0$ and, if m' denotes the number of terms in the sum, then $1 \leq m' < m$. Now repeat the argument: Either $\sum_{j \in S} \eta_j \sqrt{n_j} \neq 0$ for any choice of $\eta_j = \pm 1$, in which case Lemma 6 gives that

$$\left| \sum_{j \in S} \epsilon_j \sqrt{n_j} \right| \geq \frac{1}{(m' \sqrt{M})^{2^{m'-1}-1}} > \frac{1}{(m \sqrt{M})^{2^{m-1}-1}}$$

or else one can find a further subdivide the set S as before. Since the number of terms in the sum is a positive integer and reduces upon each subdivision, this process terminates. $\square$

Proof of Proposition 5. Expanding (7) out,

$$\begin{aligned} \mathcal{M}_m &= \frac{2^m}{\pi^m} \sum_{n_1, \dots, n_m \geq 1} \prod_{j=1}^m \frac{r(n_j)}{n_j^{3/4}} \sin\left(\frac{\pi \sqrt{n_j}}{L}\right) \widehat{\psi}\left(\sqrt{\frac{n_j}{M}}\right) \\ &\quad \times \left\langle \prod_{j=1}^m \sin\left(2\pi\left(t + \frac{1}{2L}\right)\sqrt{n_j} + \frac{1}{4}\pi\right) \right\rangle \end{aligned}$$

Now,

$$\begin{aligned} &\left\langle \prod_{j=1}^m \sin\left(2\pi\left(t + \frac{1}{2L}\right)\sqrt{n_j} + \frac{1}{4}\pi\right) \right\rangle \\ &= \left\langle \prod_{j=1}^m \frac{1}{2i} \left[e^{2\pi i((t+1/2L)\sqrt{n_j}+1/8)} - e^{-2\pi i((t+1/2L)\sqrt{n_j}+1/8)} \right] \right\rangle \\ &= \sum_{\epsilon_j = \pm 1} \frac{1}{2^m i^m} \int_{-\infty}^{\infty} \prod_{j=1}^m \epsilon_j \exp\left(\epsilon_j 2\pi i \left((t + \frac{1}{2L})\sqrt{n_j} + \frac{1}{8}\right)\right) \frac{1}{T} \omega\left(\frac{t}{T}\right) dt \\ &= \sum_{\epsilon_j = \pm 1} \frac{\prod \epsilon_j}{2^m i^m} \widehat{\omega}\left(-T \sum_{j=1}^m \epsilon_j \sqrt{n_j}\right) e^{\sum_{j=1}^m \epsilon_j \pi i (\frac{1}{T} \sqrt{n_j} + 1/4)} \end{aligned}$$

By the compact support condition of $\widehat{\psi}$, we may always assume that $n_j = \mathcal{O}(M)$. By Lemma 7 and the fact that $\widehat{\omega}$ decays faster than any polynomial power, the off-diagonal terms (those terms with $\sum_{j=1}^m \epsilon_j \sqrt{n_j} \neq 0$) contribute at most

$$\sum_{1 \leq n_1, \dots, n_m \leq M} \left(\frac{(\sqrt{M})^{2^{m-1}-1}}{T} \right)^A \ll \frac{M^{(2^{m-1}-1)A/2+m}}{T^A} \ll T^{-\delta A + 2m/(2^{m-1}-1)}$$

which is vanishingly small, since A can be arbitrarily large. Thus the only contributing terms are those with $\sum_{j=1}^m \epsilon_j \sqrt{n_j} = 0$, and using the fact that $\widehat{\omega}(0) = 1$, we therefore have for any $B > 0$,

$$\mathcal{M}_m = \sum_{n_1, \dots, n_m} \sum_{\substack{\epsilon_j = \pm 1 \\ \sum \epsilon_j \sqrt{n_j} = 0}} \prod_{j=1}^m \frac{-i\epsilon_j r(n_j)}{\pi n_j^{3/4}} \sin\left(\frac{\pi \sqrt{n_j}}{L}\right) \widehat{\psi}\left(\sqrt{\frac{n_j}{M}}\right) e^{i\pi \epsilon_j / 4} + \mathcal{O}(T^{-B})$$

In order to estimate the size of $\mathcal{M}_m/\sigma^m$ when $L \rightarrow \infty$, we need to use a lemma of Besicovitch [2].

Lemma 8. *If q_j , (for $j = 1, \dots, m$), are distinct squarefree positive integers, then $\sqrt{q_1}, \dots, \sqrt{q_m}$ are linearly independent over the rationals.*

Therefore, if $\sum_{j=1}^m \epsilon_j \sqrt{n_j} = 0$ with $n_j \geq 1$, then there must exist a division of $\{1, \dots, m\}$ into $\{S_i\}$ such that

$$\{1, \dots, m\} = \coprod_{i=1}^{\ell} S_i$$

where $\sum_{i=1}^{\ell} |S_i| = m$ such that for $i = 1, 2, \dots, \ell$, for all $j \in S_i$, $n_j = q_i f_j^2$ with the q_i being distinct square-free integers, and with the f_j satisfying

$$\sum_{j \in S_i} \epsilon_j f_j = 0$$

Summing over all possible divisions we see that

$$\begin{aligned} \frac{\mathcal{M}_m}{\sigma^m} &= \sum_{\ell=1}^m \sum_{\{1, \dots, m\} = \coprod_{i=1}^{\ell} S_i} \left(\frac{1}{\sigma^{|S_1|}} \sum_{q_1 \square \text{free}} D_{q_1}(S_1) \right) \\ &\quad \times \left(\frac{1}{\sigma^{|S_2|}} \sum_{\substack{q_2 \square \text{free} \\ q_2 \neq q_1}} D_{q_2}(S_2) \right) \cdots \left(\frac{1}{\sigma^{|S_{\ell}|}} \sum_{\substack{q_{\ell} \square \text{free} \\ q_{\ell} \neq q_1, \dots, q_{\ell-1}}} D_{q_{\ell}}(S_{\ell}) \right) \end{aligned}$$

where

$$(11) \quad D_q(S) :=$$

$$\frac{1}{q^{3|S|/4}} \sum_{\substack{f_j \geq 1 \\ \epsilon_j = \pm 1 \\ \sum_{j \in S} \epsilon_j f_j = 0}} \prod_{j \in S} \frac{-i\epsilon_j e^{i\pi \epsilon_j / 4} r(q f_j^2)}{\pi f_j^{3/2}} \sin\left(\pi \frac{1}{L} f_j \sqrt{q}\right) \widehat{\psi}\left(f_j \sqrt{\frac{q}{M}}\right)$$

We will show in Lemma 9 that if $L \rightarrow \infty$ such that $L/\sqrt{M} \rightarrow 0$ then for all $\delta' > 0$,

$$\frac{1}{\sigma^{|S|}} \sum_{q \sqsubseteq \text{free}} D_q(S) = \begin{cases} 0 & \text{if } |S| = 1 \\ 1 & \text{if } |S| = 2 \\ \mathcal{O}\left(\frac{1}{L^{1-\delta'}}\right) & \text{otherwise} \end{cases}$$

Therefore the only terms in $\mathcal{M}_m/\sigma^m$ which do not vanish as $L \rightarrow \infty$ are those where $|S_i| = 2$ for all i . If m is odd, there are no such terms, and if $m = 2k$ is even, then the number of terms is equal to the number of ways of partitioning $\{1, \dots, 2k\}$ into $\coprod_{i=1}^k S_i$ with $|S_i| = 2$, which equals

$$\frac{1}{k!} \binom{2k}{2} \binom{2k-2}{2} \cdots \binom{2}{2} = \frac{(2k)!}{k! 2^k}$$

This completes the proof of Proposition 5 □

Lemma 9. *If $L \rightarrow \infty$ is such that $L/\sqrt{M} \rightarrow 0$ then*

$$\frac{\sum_{q \sqsubseteq \text{free}} D_q(S)}{\sigma^{|S|}} = \begin{cases} 1 & \text{if } |S| = 2 \\ \mathcal{O}\left(\frac{1}{L^{1-\delta}}\right) & \text{otherwise} \end{cases}$$

for all $\delta > 0$, where $D_q(S)$ is defined in (11), and σ^2 is defined in (9).

Proof. For convenience we assume, without loss of generality, that $S = \{1, 2, \dots, |S|\}$. Using $r(n) \ll n^\delta$ for all $\delta > 0$, and $\widehat{\psi}(x) \ll 1$, we can upper bound by

$$(12) \quad \sum_{q \sqsubseteq \text{free}} D_q(S) \ll \sum_{q=1}^{\infty} \frac{q^{|S|\delta}}{q^{3|S|/4}} Q(q)$$

where

$$Q(q) = \sum_{\epsilon_j = \pm 1} \sum_{\substack{f_j \geq 1 \\ \sum_{j=1}^{|S|} \epsilon_j f_j = 0}} \prod_{j=1}^{|S|} \frac{f_j^\delta}{f_j^{3/2}} |\sin(\pi f_j \sqrt{q}/L)|$$

Note that $Q(q) \ll 1$ for all q . When $q \ll L^2$ a sharper result can be deduced by a more careful treatment of $Q(q)$. In order for $\sum_{j=1}^{|S|} \epsilon_j f_j = 0$, at least two of the ϵ must have different signs, and so, with no loss of generality, we put $\epsilon_{|S|} = -1$ and $\epsilon_{|S|-1} = +1$. Hence

$$f_{|S|} = f_{|S|-1} + \sum_{j=1}^{|S|-2} \epsilon_j f_j$$

In order for both $f_{|S|} \geq 1$ and $f_{|S|-1} \geq 1$, it must be that

$$f_{|S|-1} \geq 1 + \max \left\{ 0, - \sum_{j=1}^{|S|-2} \epsilon_j f_j \right\}$$

Therefore,

$$Q(q) = 2 \sum_{\epsilon_1, \dots, \epsilon_{|S|-2} = \pm 1} \sum_{f_1, \dots, f_{|S|-2} \geq 1} \sum_{f_{|S|-1} \geq 1 + \max\{0, -\sum_{j=1}^{|S|-2} \epsilon_j f_j\}} \left(\prod_{j=1}^{|S|-1} \frac{|\sin(\pi f_j \sqrt{q}/L)|}{f_j^{3/2-\delta}} \right) \frac{|\sin(\pi \frac{\sqrt{q}}{L} (f_{|S|-1} + \sum_{j=1}^{|S|-2} \epsilon_j f_j))|}{(f_{|S|-1} + \sum_{j=1}^{|S|-2} \epsilon_j f_j)^{3/2-\delta}}$$

Changing sums into integrals gives

$$Q(q) \ll \int \cdots \int_1^\infty dx_1 \dots dx_{|S|-2} \sum_{\epsilon_j = \pm 1} \int_{1 + \max\{0, -\sum_{j=1}^{|S|-2} \epsilon_j x_j\}}^\infty dx_{|S|-1} \times \left(\prod_{j=1}^{|S|-1} \frac{|\sin(\pi \frac{1}{L} x_j \sqrt{q})|}{x_j^{3/2-\delta}} \right) \frac{|\sin(\pi \frac{1}{L} \sqrt{q} (x_{|S|-1} + \sum_{j=1}^{|S|-2} \epsilon_j x_j))|}{(x_{|S|-1} + \sum_{j=1}^{|S|-2} \epsilon_j x_j)^{3/2-\delta}}$$

and changing variables to $x_j \sqrt{q}/L \rightarrow y_j$,

$$Q(q) \ll \frac{q^{|S|/4+1/2-|S|\delta/2}}{L^{|S|/2+1-|S|\delta}} \int \cdots \int_{\sqrt{q}/L}^\infty \sum_{\epsilon_j = \pm 1} \int_{\max\{0, -\sum \epsilon_j y_j\} + \sqrt{q}/L}^\infty \left(\prod_{j=1}^{|S|-1} \frac{|\sin(\pi y_j)|}{y_j^{3/2-\delta}} \right) \frac{|\sin(\pi (y_{|S|-1} + \sum_{j=1}^{|S|-2} \epsilon_j y_j))|}{(y_{|S|-1} + \sum_{j=1}^{|S|-2} \epsilon_j y_j)^{3/2-\delta}} dy_{|S|-1} dy_{|S|-2} \dots dy_1$$

Since the multiple integral is bounded, we may conclude that

$$Q(q) \ll \begin{cases} \frac{q^{|S|/4+1/2-|S|\delta/2}}{L^{|S|/2+1-|S|\delta}} & \text{if } q < L^2 \\ 1 & \text{if } q \geq L^2 \end{cases}$$

substituting this into (12) we see that

$$\sum_{q \square \text{free}} D_q(S) \ll \begin{cases} \frac{L^{\delta'}}{L} & \text{if } |S| = 2 \\ \frac{L^{\delta'}}{L^{|S|/2+1}} & \text{if } |S| \geq 3 \end{cases}$$

Hence

$$\frac{1}{\sigma^{|S|}} \sum_{q \square \text{free}} D_q(S) \ll \begin{cases} L^{\delta'} & \text{if } |S| = 2 \\ \frac{1}{L^{1-\delta'}} & \text{if } |S| \geq 3 \end{cases}$$

since equation (10) gives $\sigma \sim \frac{4\sqrt{\log L}}{\sqrt{L}}$ when $L \rightarrow \infty$ but $L/\sqrt{M} \rightarrow 0$. However, in the case $|S| = 2$, by the definition of $D_q(S)$ and σ^2 we see that

$$\sum_{q \square \text{free}} D_q(S) = \sigma^2.$$

This completes the proof of the lemma. $\square$

4. UNSMOOTHING

Recall that $S(t, 1/L)$ is the normalized remainder term for the number of lattice points in an annulus of inner radius t and width $1/L$. In this section we prove Theorem 1 by showing that the variance of the difference $(S(t, 1/L) - \tilde{S}_{M,L}(t))/\sigma$ vanishes and then combining this with Chebyshev's inequality to deduce a distribution theorem for $S(t, 1/L)$.

We begin with an approximation result for $N(t)$:

Lemma 10. *For any $a > 0$, $c > 1$ we have*

$$N(t) = \pi t^2 - \frac{\sqrt{t}}{\pi} \sum_{n \leq X} \frac{r(n)}{n^{3/4}} \cos(2\pi t \sqrt{n} + \frac{1}{4}\pi) + \mathcal{O}(|t|^{-1/2}) + \mathcal{O}(X^a) + \mathcal{O}\left(\frac{|t|^{2c-1}}{\sqrt{X}}\right)$$

This Lemma was already invoked by Heath-Brown in [9], with the proof being an argument similar to that which derives (12.4.4) in [17].

Lemma 11. *Suppose $L \rightarrow \infty$ as $T \rightarrow \infty$ and choose M so that $L/\sqrt{M} \rightarrow 0$ as $T \rightarrow \infty$ but $M = \mathcal{O}(T^{2(1-\delta)})$ (for a fixed $\delta > 0$). Then as $T \rightarrow \infty$,*

$$\left\langle |S(t, 1/L) - \tilde{S}_{M,L}(t)|^2 \right\rangle \ll \frac{\log M}{\sqrt{M}}$$

Proof. Putting $a = \delta'$ and $c = 1 + \delta'/2$ for $\delta' > 0$ arbitrarily small in Lemma 10, we have

$$\begin{aligned} S(t, 1/L) &:= \frac{N(t + 1/L) - N(t) - \pi(2t/L + 1/L^2)}{\sqrt{t}} \\ &= \frac{2}{\pi} \sum_{n \leq X} \frac{r(n)}{n^{3/4}} \sin\left(\frac{\pi\sqrt{n}}{L}\right) \sin\left(2\pi\left(t + \frac{1}{2L}\right)\sqrt{n} + \frac{1}{4}\pi\right) + R(X, t) \end{aligned}$$

where

$$R(X, t) \ll \frac{1}{|t|} + \frac{X^{\delta'}}{\sqrt{|t|}} + \frac{|t|^{1/2+\delta'}}{\sqrt{X}}$$

Set $X = T^{2-\delta}$. Since $M = \mathcal{O}(T^{2(1-\delta)})$ and $\hat{\psi}$ has compact support, the infinite sum in $\tilde{S}_{M,L}(t)$, given in (6), is truncated before $n = T^{2-\delta}$, and so

$$\begin{aligned} S(t, 1/L) - \tilde{S}_{M,L}(t) &= \\ \frac{2}{\pi} \sum_{n \leq T^{2-\delta}} \frac{r(n)}{n^{3/4}} \sin\left(\frac{\pi\sqrt{n}}{L}\right) \sin\left(2\pi\left(t + \frac{1}{2L}\right)\sqrt{n} + \frac{\pi}{4}\right) &\left(1 - \hat{\psi}\left(\sqrt{\frac{n}{M}}\right)\right) + R(T^{2-\delta}, t) \end{aligned}$$

Let P denote the sum, then Cauchy-Schwartz gives

$$(13) \quad \left\langle (S(t, 1/L) - \tilde{S}_{M,L}(t))^2 \right\rangle = \langle P^2 \rangle + \langle R(T^{2-\delta}, t)^2 \rangle + \mathcal{O}\left(\sqrt{\langle P^2 \rangle} \sqrt{\langle R(T^{2-\delta}, t)^2 \rangle}\right)$$

Observe that

$$\langle R(T^{2-\delta}, t)^2 \rangle \ll T^{-1+\delta''}$$

for arbitrarily small $\delta'' > 0$, and

$$\begin{aligned} \langle P^2 \rangle &= \frac{2}{\pi^2} \sum_{n \leq T^{2-\delta}} \frac{r(n)^2}{n^{3/2}} \sin^2 \left(\frac{\pi \sqrt{n}}{L} \right) \left(1 - \widehat{\psi} \left(\sqrt{\frac{n}{M}} \right) \right)^2 \\ &\quad + \mathcal{O} \left(\sum_{1 \leq m \neq n \leq T^{2-\delta}} \widehat{\omega}(T(\sqrt{n} - \sqrt{m})) \right) \end{aligned}$$

The same argument used in §3.1 shows the error term here vanishes like $\mathcal{O}(T^{-B})$ for any $B > 0$.

Since $\sum_{n \leq X} r(n)^2 \sim 4X \log X$, partial summation gives

$$\begin{aligned} \langle P^2 \rangle &\sim \frac{8}{\pi^2} \int_1^{T^{2-\delta}} \frac{\log x}{x^{3/2}} \sin^2 \left(\frac{\pi \sqrt{x}}{L} \right) \left(1 - \widehat{\psi} \left(\sqrt{\frac{x}{M}} \right) \right)^2 dx \\ &= \frac{32}{L\pi^2} \int_{1/L}^{T/L} \frac{\log(Ly) \sin^2(\pi y)}{y^2} \left(1 - \widehat{\psi} \left(\frac{yL}{\sqrt{M}} \right) \right)^2 dy \end{aligned}$$

by change of variables $x = y^2 L^2$. If $yL/\sqrt{M} \ll 1$ then

$$\widehat{\psi} \left(\frac{yL}{\sqrt{M}} \right) = 1 + \mathcal{O} \left(\frac{yL}{\sqrt{M}} \right)$$

leading to

$$\begin{aligned} \langle P^2 \rangle &\ll \frac{L}{M} \int_0^{\sqrt{M}/L} \log(Ly) \sin^2(\pi y) dy + \frac{1}{L} \int_{\sqrt{M}/L}^{T/L} \frac{\log(Ly) \sin^2(\pi y)}{y^2} dy \\ &\ll \frac{\log M}{\sqrt{M}} \end{aligned}$$

Inserting this into (13), using $M = \mathcal{O}(T^{2(1-\delta)})$ and choosing $0 < \delta'' < \delta$ in the estimate of $\langle R(X, t)^2 \rangle$ we have that

$$\begin{aligned} \left\langle (S(t, 1/L) - \widetilde{S}_{M,L}(t))^2 \right\rangle &\ll \frac{\log M}{\sqrt{M}} + \frac{1}{T^{1-\delta''}} + \frac{\sqrt{\log M}}{M^{1/4} T^{1/2-\delta''/2}} \\ &= \mathcal{O} \left(\frac{\log M}{\sqrt{M}} \right) \end{aligned}$$

□

Lemma 12. *Under the conditions of Lemma 11, we have for all fixed $\eta > 0$,*

$$\mathbb{P}_{\omega, T} \left\{ \left| \frac{S(t, 1/L)}{\sigma} - \frac{\widetilde{S}_{M,L}(t)}{\sigma} \right| > \eta \right\} \rightarrow 0$$

as $T \rightarrow \infty$, where $\sigma^2 = 16 \log L/L$.

Proof. For fixed $\eta > 0$, Chebychev's inequality gives

$$\begin{aligned} \mathbb{P}_{\omega, T} \left\{ \left| \frac{S(t, 1/L)}{\sigma} - \frac{\widetilde{S}_{M,L}(t)}{\sigma} \right| > \eta \right\} &\leq \frac{\left\langle (S(t, 1/L) - \widetilde{S}_{M,L}(t))^2 \right\rangle}{\eta^2 \sigma^2} \\ &\ll \frac{L}{\log L} \frac{\log M}{\sqrt{M}} \end{aligned}$$

which tends to zero as $T \rightarrow \infty$ by the assumptions placed on M and L .

□

Corollary 13. *If $L \rightarrow \infty$ but $L = \mathcal{O}(T^\delta)$ for all $\delta > 0$ as $T \rightarrow \infty$, then for any interval $\mathcal{A}$,*

$$(14) \quad \mathbb{P}_{\omega, T} \left\{ \frac{S(t, 1/L)}{\sigma} \in \mathcal{A} \right\} \rightarrow \frac{1}{\sqrt{2\pi}} \int_{\mathcal{A}} e^{-x^2/2} dx$$

where $\sigma^2 = 16 \log L/L$.

Proof. Set $M = L^3$, then $M = \mathcal{O}(T^\delta)$ for all $\delta > 0$ and $L/\sqrt{M} \rightarrow 0$. Thus $\tilde{S}_{M,L}/\sigma$ weakly converges to a standard normal distribution as $T \rightarrow \infty$ when t is smoothly averaged around T by Theorem 2. But Lemma 12 implies that $S(t, 1/L)/\sigma$ must also weakly converge to a standard normal distribution too. $\square$

We are now able to prove our main result, Theorem 1, which says that if $L \rightarrow \infty$ but $L = \mathcal{O}(T^\delta)$ for all $\delta > 0$, then for any interval $\mathcal{A}$

$$\lim_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ t \in [T, 2T] : \frac{S(t, 1/L)}{\sigma} \in \mathcal{A} \right\} = \frac{1}{\sqrt{2\pi}} \int_{\mathcal{A}} e^{-x^2/2} dx$$

Proof of Theorem 1. Fix $\epsilon > 0$, and approximate the indicator function $\mathbb{1}_{[1,2]}$ above and below by smooth functions $\chi_{\pm} \geq 0$ so that $\chi_- \leq \mathbb{1}_{[1,2]} \leq \chi_+$, where both $\chi_{\pm}$ and their Fourier transforms are smooth and of rapid decay, and so that their total masses are within ϵ of unity: $|\int \chi_{\pm}(x) dx - 1| < \epsilon$. Now set $\omega_{\pm} := \chi_{\pm}/\int \chi_{\pm}$. Then $\omega_{\pm}$ are “admissible” and for all t ,

$$(15) \quad (1 - \epsilon)\omega_-(t) \leq \mathbb{1}_{[1,2]}(t) \leq (1 + \epsilon)\omega_+(t)$$

Now

$$\text{meas} \left\{ t \in [T, 2T] : \frac{S(t, 1/L)}{\sigma} \in \mathcal{A} \right\} = \int_{-\infty}^{\infty} \mathbb{1}_{\mathcal{A}} \left(\frac{S(t, 1/L)}{\sigma} \right) \mathbb{1}_{[1,2]} \left(\frac{t}{T} \right) dt$$

and since (15) holds, we find

$$\begin{aligned} (1 - \epsilon)\mathbb{P}_{\omega_-, T} \left\{ \frac{S(t, 1/L)}{\sigma} \in \mathcal{A} \right\} &\leq \frac{1}{T} \text{meas} \left\{ t \in [T, 2T] : \frac{S(t, 1/L)}{\sigma} \in \mathcal{A} \right\} \\ &\leq (1 + \epsilon)\mathbb{P}_{\omega_+, T} \left\{ \frac{S(t, 1/L)}{\sigma} \in \mathcal{A} \right\} \end{aligned}$$

By Corollary 13 the two extreme sides of this inequality have a limit as $T \rightarrow \infty$, of

$$(1 \pm \epsilon) \frac{1}{\sqrt{2\pi}} \int_{\mathcal{A}} e^{-x^2/2} dx$$

and so we get that

$$(1 - \epsilon) \frac{1}{\sqrt{2\pi}} \int_{\mathcal{A}} e^{-x^2/2} dx \leq \liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ t \in [T, 2T] : \frac{S(t, 1/L)}{\sigma} \in \mathcal{A} \right\}$$

with a similar statement for limsup; since $\epsilon > 0$ is arbitrary this shows that the limit exists and equals

$$\lim_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ t \in [T, 2T] : \frac{S(t, 1/L)}{\sigma} \in \mathcal{A} \right\} = \frac{1}{\sqrt{2\pi}} \int_{\mathcal{A}} e^{-x^2/2} dx$$

which is the Gaussian law. $\square$

ACKNOWLEDGMENTS

We would like to thank Jens Marklof for helpful discussions. This work was supported in part by the EC TMR network *Mathematical Aspects of Quantum Chaos*, EC-contract no HPRN-CT-2000-00103 and the Israel Science Foundation founded by the Israel Academy of Sciences and Humanities.

REFERENCES

- [1] M.V. Berry and M. Tabor, "Level clustering in the regular spectrum", *Proc. Roy. Soc. A* **356** (1977) 375–394.
- [2] A.S. Besicovitch, "On the linear independence of fractional powers of integers", *J. London Math. Soc.* **15** (1940) 3–6.
- [3] P. Bleher, "Trace formula for quantum integrable systems, lattice-point problems and small divisors", *Emerging Applications of Number Theory*, D.A. Hejhal, J. Friedman, M.C. Gutzwiller, A.M. Odlyzko, eds. (Springer, 1999) pp. 1–38.
- [4] P. Bleher, Zh. Cheng, F. Dyson and J. Lebowitz, "Distribution for the error term for the number of lattice points inside a shifted circle", *Comm. Math. Phys.* **154** (1993) 433–469
- [5] P. Bleher, F. Dyson, and J. Lebowitz, "Non-Gaussian energy level statistics for some integrable systems". *Phys. Rev. Lett.* **71** (1993) 3047–3050.
- [6] P. Bleher and J. Lebowitz, "Energy-level statistics of model quantum systems: universality and scaling in a lattice-point problem" *J. Statist. Phys.* **74** (1994) 167–217.
- [7] P. Bleher and J. Lebowitz, "Variance of number of lattice points in random narrow elliptic strip", *Ann. Inst. H. Poincaré Probab. Statist.* **31** (1995) 27–58.
- [8] A. Eskin, G. Margulis and S. Mozes "Quadratic forms of signature (2, 2) and eigenvalue spacings on flat tori", preprint.
- [9] D.R. Heath-Brown, "The distribution and moments of the error term in the Dirichlet divisor problem" *Acta Arithmetica* **60** (1992) 389–415.
- [10] D.R. Heath-Brown, "The Dirichlet divisor problem", *Advances in Number Theory* (Kingston, ON, 1991), Oxford Sci. Publ., 1993, pp. 31–35,
- [11] P. Major, "Poisson law for the number of lattice points in a random strip with finite area". *Prob. Th. Rel. Fields* **92** (1992) 423–464.
- [12] J. Marklof, "Pair correlation densities of inhomogeneous quadratic forms", *Ann. of Math.* (2), to appear.
- [13] N. Minami, "On the Poisson limit theorems of Sinai and Major". *Comm. Math. Phys.* **213** (2000) 203–247.
- [14] S. Ramanujan "Some formulae in the analytic theory of numbers", *Messenger of Math.* **XLV** (1916) 81–84.
- [15] P. Sarnak "Values at integers of binary quadratic forms", *Harmonic Analysis and Number Theory* (Montreal, 1996), CMS Conf. Proc. **21**, 1997, pp. 181–203.
- [16] Ya. G. Sinai, "Poisson distribution in a geometric problem", *Dynamical Systems and Statistical Mechanics* (Moscow, 1991), Adv. Soviet Math., **3**, 1991, pp. 199–214.
- [17] E.C. Titchmarsh, *The Theory of the Riemann Zeta-Function* (2nd edition, revised by D.R. Heath-Brown), (Oxford Science Publications, 1986).

RAYMOND AND BEVERLY SACKLER SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, TEL AVIV 69978, ISRAEL.

Current address: American Institute of Mathematics, 360 Portage Avenue, Palo Alto, CA 94306-2244, U.S.A.

E-mail address: `hughes@aimath.org`

RAYMOND AND BEVERLY SACKLER SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, TEL AVIV 69978, ISRAEL

E-mail address: `rudnick@post.tau.ac.il`